

# LIST OF REFERENCES CITED BY APPLICANT

(Use several sheets if necessary)

ATTY. DOCKET NO.

7853-238

APPLICATION NO.

TBA

APPLICANT


Kingsbury et al.

FILING DATE

Herewith

GROUP

1644



## U.S. PATENT DOCUMENTS

| EXAMINER<br>INITIAL |    | DOCUMENT NUMBER | DATE     | NAME             | CLASS | SUBCLASS | FILING DATE<br>IF<br>APPROPRIATE |
|---------------------|----|-----------------|----------|------------------|-------|----------|----------------------------------|
| ay                  | AA | 5,840,691       | 11/24/98 | Furcht et al.    |       |          |                                  |
| ay                  | AB | 5,721,351       | 2/24/98  | Levinson         |       |          |                                  |
| ay                  | AC | 5,591,719       | 1/7/97   | Furcht et al.    |       |          |                                  |
| ay                  | AD | 5,262,311       | 11/16/93 | Pardee and Liang |       |          |                                  |
| ay                  | AE | 5,116,964       | 5/26/92  | Capon and Lasky  |       |          |                                  |

## FOREIGN PATENT DOCUMENTS

|  |    | DOCUMENT NUMBER | DATE     | COUNTRY | CLASS | SUBCLASS | TRANSLATION |    |
|--|----|-----------------|----------|---------|-------|----------|-------------|----|
|  |    |                 |          |         |       |          | YES         | NO |
|  | AF | WO 99/15663     | 4/1/99   | PCT     |       |          |             |    |
|  | AG | WO 97/42224     | 11/13/97 | PCT     |       |          |             |    |
|  | AH | JP 6-178687     | 6/28/94  | Japan   |       |          |             |    |
|  | AI | EP 239400       | 8/3/94   | EPO     |       |          |             |    |




## OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.)

|  |    |   |
|--|----|---|
|  | AJ | GenBank Accession No. E07714 (cDNA encoding mouse ST2 protein)  |
|  | AK | GenBank Accession No. P14719 (mouse ST2 protein precursor)  |
|  | AL | GenBank Accession No. E08652 (cDNA encoding mouse ST2L)   |
|  | AM | GenBank Accession No. S29498 (mouse ST2L protein)   |
|  | AN | GenBank Accession No. AB012701 (human mRNA for ST2L, 5'noncoding region incl. exon 1a, complete cds)  |
|  | AO | GenBank Accession No. BAA82405 (human ST2L protein)   |
|  | AP | GenBank Accession No. D12763 (human mRNA for ST2L protein)  |
|  | AQ | GenBank Accession No. BAA0223 (human ST2 protein)   |
|  | AR | GenBank Accession No. AB029084 (human ST2V mRNA, complete cds)  |
|  | AS | GenBank Accession No. BAA85894 (human ST2V)   |
|  | AT | GenBank Accession No. Y07519 (mouse St2 gene)   |
|  | AU | GenBank Accession No. M28621 (mouse interferon-γ mRNA sequence)   |
|  | AV | Altschul et al., 1997, "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucl Acids Res 25:3389-3402      |
|  | AW | Altschul et al., 1990, "Basic Local Alignment Search Tool", J. Mol. Biol. 215:403-410   |
|  | AX | Anderson and Coyle, 1994, "TH2 and 'TH2-like' cells in allergy and asthma: pharmacological perspectives", Trends Pharmacol. Sci. 15:324-332 |

|    |   |
|----|---|
| AY | Aruffo et al., 1990, "CD44 is the principal cell surface receptor for hyaluronate", <i>Cell</i> <b>61</b> :1303-1313  |
| AZ | Bergers et al., 1994, "Alternative promoter usage of the Fos-responsive gene <i>Fit-1</i> generates mRNA isoforms coding for either secreted or membrane-bound proteins related to the IL-1 receptor", <i>EMBO J.</i> <b>13</b> :1176-1188  |
| BA | Chen et al., 1993, "RAG-2-deficient blastocyst complementation: an assay of gene function in lymphocyte development", <i>Proc. Natl. Acad. Sci. USA</i> <b>90</b> :4528-4532  |
| BB | Clerici et al., 1993, "Restoration of HIV-specific cell-mediated immune responses by interleukin-12 in vitro", <i>Science</i> <b>262</b> :1721-1724   |
| BC | Clerici et al., 1993, "Changes in interleukin-2 and interleukin-4 production in asymptomatic, human immunodeficiency virus-seropositive individuals", <i>J. Clin. Invest.</i> <b>91</b> :759-765  |
| BD | Cohn et al., 1997, "Induction of airway mucus production by T helper 2 (Th2) cells: a critical role for interleukin 4 in cell recruitment but not mucus production", <i>J. Exp. Med.</i> <b>186</b> :1737-1747  |
| BE | Del Prete et al., 1991, "Purified protein derivative of <i>Mycobacterium tuberculosis</i> and excretory-secretory antigen(s) of <i>Toxocara canis</i> expand in vitro human T cells with stable and opposite (Type 1 T helper or Type 2 T helper) profile of cytokine production", <i>J. Clin. Invest.</i> <b>88</b> :346-350 |
| BF | Fägerstam et al., 1992, "Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis", <i>J. Chromatog.</i> <b>597</b> :397-410  |
| BG | Firestein et al., 1989, "A new murine CD4 <sup>+</sup> T cell subset with an unrestricted cytokine profile", <i>J. Immunol.</i> <b>143</b> :518-525   |
| BH | Frangogiannis et al., 1998, "Resident cardiac mast cells degranulate and release preformed TNF- $\alpha$ initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion", <i>Circulation</i> <b>98</b> :699-710   |
| BI | Gächter et al., 1996, "Transcription of the interleukin-1 receptor-related T1 gene is initiated at different promoters in mast cells and fibroblasts", <i>J. Biol. Chem.</i> <b>271</b> :124-129  |
| BJ | Gavett et al., 1994, "Depletion of murine CD4 <sup>+</sup> T lymphocytes prevents antigen-induced airway hyperreactivity and pulmonary eosinophilia", <i>Am. J. Respir Cell Mol. Biol.</i> <b>10</b> :587-593   |
| BK | George et al., 1994, "Embryonic expression and cloning of the murine GATA-3 gene", <i>Development</i> <b>120</b> :2673-2686   |
| BL | Gonazlo et al., 1996, "Mouse eotaxin expression parallels eosinophil accumulation during lung allergic inflammation but it is not restricted to a Th2-type response", <i>Immunity</i> <b>4</b> :1-14  |
| BM | Gu et al., 1994, "Deletion of a DNA polymerase $\beta$ gene segment in T cells using cell type-specific gene targeting", <i>Science</i> <b>265</b> :103-106   |
| BN | Hamelmann et al., 1997, "Noninvasive measurement of airway responsiveness in allergic mice using barometric plethysmography", <i>Am. J. Respir. Crit. Care Med.</i> <b>156</b> :766-775   |
| BO | Holgate, 1997, "The cellular and mediator basis of asthma in relation to natural history", <i>Lancet</i> <b>350</b> (Suppl. 2):5-9  |
| BP | Kanagawa et al., 1993, "Resistance of mice deficient in IL-4 to retrovirus-induced immunodeficiency syndrome (MAIDS)", <i>Science</i> <b>262</b> :240-242   |
| BQ | Kaneshima et al., 1994, "Human hematolymphoid cells in SCID mice", <i>Curr. Opin. Immunol.</i> <b>6</b> :327-333  |

|    |   |
|----|---|
| BR | Karlin and Altschul, 1993, "Applications and statistics for multiple high-scoring segments in molecular sequences", Proc. Natl. Acad. Sci. USA <u>90</u> :5873-5877   |
| BS | Karlin and Altschul, 1990, "Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes", Proc Natl Acad Sci USA <u>87</u> :2264-2268  |
| BT | Klemenz et al., 1989, "Serum- and oncoprotein-mediated induction of a gene with sequence similarity to the gene encoding carcinoembryonic antigen", Proc. Natl. Acad. Sci. USA <u>86</u> :5708-5712   |
| BU | Ko and Engel, 1993, "DNA binding specificities of the GATA transcription factor family", Mol. Cell. Biol. <u>13</u> :4011-4022  |
| BV | Kumar et al., 1997, "Expression of ST2, an interleukin-1 receptor homologue, is induced by proinflammatory stimuli", Biochem. Biophys. Res. Comm. <u>235</u> :474-478   |
| BW | Liew, 1994, "Induction and regulation of CD4 <sup>+</sup> T cell subsets", CIBA Foundation Symposium <u>87</u> :170-178   |
| BX | Löhning et al., 1998, "T1/ST2 is preferentially expressed on murine Th2 cells, independent of interleukin 4, interleukin 5, and interleukin 10, and important for Th2 effector function", Proc. Natl. Acad. Sci. USA <u>95</u> :6930-6935                           |
| BY | Lukacs et al., 1994, "Interleukin-4-dependent pulmonary eosinophil infiltration in a murine model of asthma", Am. J. Respir. Cell Mol. Biol. <u>10</u> :526-532   |
| BZ | Maggi et al., 1994, "Th2-like CD8 <sup>+</sup> T cells showing B cell helper function and reduced cytolytic activity in human immunodeficiency virus type 1 infection", J. Exp. Med. <u>180</u> :489-495  |
| CA | Maggi et al., 1994, "Ability of HIV to promote a T <sub>H</sub> 1 to T <sub>H</sub> 0 shift and to replicate preferentially in T <sub>H</sub> 2 to T <sub>H</sub> 0 cells", Science <u>265</u> :244-248   |
| CB | Makino et al., 1990, "H-2-associated and background genes influence the development of a murine retrovirus-induced immunodeficiency syndrome", J. Immunol. <u>144</u> :4347-4355  |
| CC | Manetti et al., 1994, "CD30 expression by CD8 <sup>+</sup> T cells producing type 2 helper cytokines. Evidence for large numbers of CD8 <sup>+</sup> CD30 <sup>+</sup> T cell clones in human immunodeficiency virus infection", J. Exp. Med. <u>180</u> :2407-2411 |
| CD | McMahan et al., 1991, "A novel IL-1 receptor, cloned from B cells by mammalian expression, is expressed in many cell types", EMBO J. <u>10</u> :2821-2832   |
| CE | Metzler and Xu, 1997, "The role of mast cells in atherosclerosis", Int. Arch. Allergy Immunol. <u>114</u> :10-14  |
| CF | Moll et al., 1991, "Expression of T-cell-associated serine proteinase 1 during murine <i>Leishmania major</i> infection correlates with susceptibility to disease", Infect. Immun. <u>59</u> :4701-4705   |
| CG | Mosmann and Moore, 1991, "The role of IL-10 in crossregulation of T <sub>H</sub> 1 and T <sub>H</sub> 2 responses", Immunol Today <u>12</u> :A49-A53  |
| CH | Mosmann and Coffman, 1989, "TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties", Annu. Rev. Immunol. <u>7</u> :145-173   |
| CI | Murphy et al., 1990, "Induction by antigen of intrathymic apoptosis of CD4 <sup>+</sup> CD8 <sup>+</sup> TCR <sup>b</sup> thymocytes in vivo", Science <u>250</u> :1720-1723  |
| CJ | Myers and Miller, 1988, "Optimal Alignments in Linear Space", Comput. Appl. Biosci. <u>4</u> :11-17   |

|    |  |
|----|--|
| CK | Oettgen and Ghea, 1999, "IgE in asthma and atopy: cellular and molecular connections", J. Clin. Invest. <u>104</u> :829-835  |
| CL | Onashi et al., 1992, "Airway hyperresponsiveness, increased intracellular spaces of bronchial epithelium, and increased infiltration of eosinophils and lymphocytes in bronchial mucosa in asthma", Am. Rev Respir. Dis. <u>145</u> :1469-1476   |
| CM | Ray and Cohn, 1999, "Th2 cells and GATA-3 in asthma: new insights into the regulation of airway inflammation", J. Clin. Invest. <u>104</u> :985-993  |
| CN | Robinson et al., 1993, "Activation of CD4 <sup>+</sup> T cells, increased T <sub>H2</sub> -type cytokine mRNA expression, eosinophil recruitment in bronchoalveolar lavage after allergen inhalation challenge in patients with atopic asthma", J. Allergy Clin. Immunol. <u>92</u> :313-324 |
| CO | Romagnani, 1997, "The Th1/Th2 paradigm", Immunol. Today <u>18</u> :263-266   |
| CP | Schweitzer and Sharpe, 1998, "Studies using antigen-presenting cells lacking expression of both B7-1 (CD80) and B7-2 (CD86) show distinct requirements for B7 molecules during priming versus restimulation of Th2 but not Th1 cytokine production", J. Immunol. <u>161</u> :2762-2771       |
| CQ | Seder et al., 1994, "CD28-mediated costimulation of interleukin 2 (IL-2) production plays a critical role in T cell priming for IL-4 and interferon- $\gamma$ production", J. Exp. Med. <u>179</u> :299-304  |
| CR | Seder and Gros, 1995, "The functional role of CD8 <sup>+</sup> T helper type 2 cells", J. Exp. Med. <u>181</u> :5-7  |
| CS | Shinkai et al., 1992, "RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement", Cell <u>68</u> :855-867   |
| CT | Tepper et al., 1990, "IL-4 induces allergic-like inflammatory disease and alters T cell development in transgenic mice", Cell <u>62</u> :457-467   |
| CU | Tominaga et al., 1999, "Presence and expression of a novel variant form of ST2 gene product in human leukemic cell line UT-7/GM", Biochem. Biophys. Res. Comm. <u>264</u> :14-18   |
| CV | Tominaga et al., 1992, "Nucleotide sequence of a complementary DNA for human ST2", Biochim. Biophys. Acta <u>1171</u> :215-218   |
| CW | Tominaga et al., 1991, "Molecular cloning of the murine ST2 gene. Characterization and chromosomal mapping", Biochim. Biophys. Acta <u>1090</u> :1-8   |
| CX | Tominaga, 1989, "A putative protein of a growth specific cDNA from BALB/c-3T3 cells is highly similar to the extracellular portion of mouse interleukin 1 receptor", FEBS Lett. <u>258</u> :301-304  |
| CY | Tsuyuki et al., 1997, "Costimulation through B7-2 (CD86) is required for the induction of a lung mucosal T helper cell 2 (Th2) immune response and altered airway responsiveness", J. Exp. Med. <u>185</u> :1671-1679  |
| CZ | Volmer et al., 1994, "T lymphocytes derived from skin lesions of patients with psoriasis vulgaris express a novel cytokine pattern that is distinct from that of T helper type 2 cells", Eur. J. Immunol. <u>24</u> :2377-2382   |
| DA | Wang et al., 1992, "IL-4 activates a distinct signal transduction cascade from IL-3 in factor-dependent myeloid cells", EMBO J. <u>11</u> :4899-4908   |
| DB | Werenskiold, 1992, "Characterization of a secreted glycoprotein of the immunoglobulin superfamily inducible by mitogen and oncogene", Eur. J. Biochem. <u>204</u> :1041-1047   |
| DC | Werenskiold et al., 1989, "Induction of a mitogen-responsive gene after expression of the Ha-ras oncogene in NIH 3T3 fibroblasts", Mol. Cell. Biol. <u>9</u> :5207-5214  |

|  |   |          |                 |   |         |
|--|---|----------|-----------------|---|---------|
| DD   | Wierenga et al., 1990, "Evidence for compartmentalization of functional subsets of CD4 <sup>+</sup> T lymphocytes in atopic patients", J. Immunol. <u>144</u> :4651-4656        |          |                 |   |         |
| DE   | Yamamura et al., 1991, "Defining protective responses to pathogens: Cytokine profiles in leprosy lesions", Science <u>254</u> :277-279  |          |                 |   |         |
| DF   | Yanagisawa et al., 1993, "Presence of a novel primary response gene ST2L, encoding a product highly similar to the interleukin 1 receptor type 1", FEBS Lett. <u>318</u> :83-87 |          |                 |   |         |
| DG   | Young et al., 1992, "Expression of cytolytic mediators by synovial fluid lymphocytes in rheumatoid arthritis", Am. J. Path. <u>140</u> :1261-1268                               |          |                 |   |         |
| DH   | Zheng and Flavell, 1997, "The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells", Cell <u>89</u> :587-596                 |          |                 |   |         |
| DI   | GenBank Accession No. E07716, cDNA encoding human ST2, published June 1994  |          |                 |   |         |
| <table border="1"> <tr> <td>EXAMINER</td> <td>DATE CONSIDERED</td> </tr> <tr> <td></td> <td>9-23-05</td> </tr> </table>   |   | EXAMINER | DATE CONSIDERED |  | 9-23-05 |
| EXAMINER   | DATE CONSIDERED   |          |                 |   |         |
|   | 9-23-05   |          |                 |   |         |
| <p>*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.</p> |   |          |                 |   |         |